

“Impact of Nitrogen and Sulphur on growth, yield and quality of Mustard [Brassica Juncea (L.) Czern and Coss] varieties”

Nirmal Chaudhary

Research Scholar

Department of Botany

Sainath University, Ormanjhi. Ranchi, Jharkhand.

Dr. S.H.A. Kazmi

Associate Prof.

Shia P.G.College, Department of Botany, Lucknow.

ABSTRACT

An experiment entitled, "Effect of nitrogen and sulphur on growth, yield and quality of mustard [Brassica Juncea (L.) Czern and Coss] varieties was conducted during rabi seasons of 2016-17 and 2018-19. The experiment was conducted in a split plot design with three replications taking three levels of nitrogen (0, 50 and 100 kg N ha⁻¹) three sulphur levels (0, 25 and 50 kg S ha⁻¹) and three mustard varieties (Vardan, Bio 902 and RH 30). The soil of the field was low in nitrogen and medium in phosphorus and potassium and slightly alkaline in reaction.

1.1 INTRODUCTION

Oilseeds occupy an important place in Indian economy as well as in human life. They are not only rich sources of energy and carriers of fat soluble vitamins A, D, E, and K but are major sources of raw materials for a wide range of industrial products also. More than 80 per cent of our country's requirements of vegetable oils and fats are derived from seven annual oilseeds crops viz. ground nut, rapeseed mustard, sesamum, safflower, sunflower, niger, soybean and two non-edible oilseeds, linseed and castor.

The average annual production of oilseeds in four years (2013-17) after launching of Technology Mission on Oilseeds, had increased from 12.0 to 16.7 million tonnes, a record annual 10 percent increase. To meet the edible oil requirement of the burgeoning population, the present level of oilseeds production (21.3 million tonnes in the year 2016-17-Economic Survey) will have to be boosted to 26.0 million tonnes by the end of this century.

The rapeseed and mustard is the second important oilseed crop, contributed about 29 percent of total oil seed production in the country during 2014-15. Area under this group of crops is spread over the states of Uttar Pradesh, Rajasthan, Punjab, Haryana, Madhya Pradesh, Bihar, Gujarat, West Bengal and Assam. However, its productivity remains low, because it is grown mostly as rainfed crop under low fertility conditions with traditional varieties. To enhance the productivity, we should bring more area under irrigation, use of balanced fertilizers and introduce newly developed promising varieties.

Among the nutrients, nitrogen and sulphur have important role in seed protein and oil synthesis. Work so far done indicated positive role of both of these nutrients in promoting yield and quality of seed of mustard. However, the quantum effect varies depending upon agro-ecosystem occupied by the crop. Among the primary

nutrients, these crops have shown higher response to nitrogen fertilization compared to phosphorus (Gangasaran and Giri, 2015). Response to nitrogen upto 80-100 kg/ha has been reported by Chahal et al. 2010. Sulphur deficiency of late has been widely recognised under light textured soils and especially in areas where intensive agriculture is done. A large number of alluvial soils where mustard is mostly cultivated, contains less than 10 ppm available S (Naik and Das, 2014). Oleiferous brassicae in general have high sulphur requirement owing to high sulphur containing amino acids in them. Sulphur is a key element in rapeseed-mustard oil and its deficiency causes a significant reduction in the oil content (Herath and Ormrod, 2011). The combined effect of nitrogen and sulphur on brassicae yield (Vaidya et al., 2011 and Tiwari, 2015) and quality (Aulakh et al. 2010) have been reported. However the work so far done has been insufficient and calls for concerted efforts to improve of this crop.

Recently a number of varieties have been developed by different centres but their performance varies from place to place depending upon soil and climate. These varieties also reacts to various inputs including nitrogen and sulphur applications. Much information is not available on these aspects especially under western parts of Uttar Pradesh.

The reproductive growth is related to vegetative growth of a plant which mainly includes the height, number of primary, secondary and total number of branches, varieties differed in respect of yield attributes viz. number of siliquae per plant, length of siliqua, number of grains/siliqua, weight of grains/plant and 1000 grain weight. Variety Vardan proved significantly superior than Bio-902 and RH-30 in respect of these yield attributing characters except 1000 grain weight where variety RH-30 produced bolder grains which in turn gave higher 1000 grain weight. Differences in yield attributing characters due to varieties have also been reported by Saik and Bhargava (2005), Sharma et al (2009), Garnayak et al (2000), Singh et al (2000), Kumar et al (2001) and Singh et al (2001).

The seed yield of mustard was greatly influenced by varieties. Variety Vardan produced significantly higher grain yield during both years. The higher seed yield of variety Vardan may be attributed to higher number of siliquae/plant, number of grains/siliqua and weight of grains per plant in comparison to Bio-902 and RH-30. Variety RH-30 though produced bolder grains and consequently higher 1000 grain weight but this could not compensate the higher number of siliquae/plant, grains/siliqua and total grain weight/plant of variety Vardan. Difference in grain yield of mustard due to varieties have also been reported by Bhan (2001), Kumar and Malik (2003), Saran and Giri (2004), Kumar and Singh (2005), Shyamraj (2006), Kumar and Singh (2009), Sharma et al (2009), Singh (2009), Singh et al (2016), Singh and Rajput (2016), Garnayak et al (2000), Singh et al (2000), Kumar et al (2001) and Singh et al (2001).

Stover yield is the result of alround development of a plant. Variety Vardan having higher number of branches produced higher stover yield. Harvest index remained unaffected by varieties during both years of study. Saik and Bhargava (2005), Sharma et al (2009), Singh et al (2016) and Kumar et al (2001), however, recorded significant differences in harvest index of mustard varieties in their studies.

Variety Vardan produced significantly higher total dry matter per hectare. This may be attributed to higher grain and stover yield of variety Vardan as compared to variety Bio-902 and RH-30.

Nitrogen content (%) in seed and stover and nitrogen uptake differed significantly due to varieties. The N content was higher in seeds and stover of variety Vardan than Bio-902 and RH-30 during both years of study. Since uptake is related to its content in seed and stover and the grain and stover yield of the plant, variety Vardan consequently recorded higher uptake of N. Varietal differences in N uptake have also been observed by Bishnoi and Singh (2000).

Sulphur content and uptake was higher due to variety Vardan in comparison to Bio-902 and RH-30. The higher uptake of S in this variety is also because of its higher grain and stover yield.

Among the qualities, oil content (%) is the most important, oil content is influenced by external characters to some extent and varieties as well being genetically different. Variety Vardan had higher oil content in its grain than Bio-902 and RH-30. The oil yield was also higher in Vardan because of higher grain

yield of this cultivar. Differences in oil content and oil yield due to varieties have also been reported by Kumar and Rahman (1998), Rastogi (2000), Bhan (2001), Saran and De (2009), Bishnoi and Singh (2009) and Singh et al (2016).

Protein content and protein yield differed due to varieties. Variety RH-30 had higher protein content in its seeds but the protein yield remained significantly lower as compared to Vardan and Bio 902. This may be due to its lowest grain yield ha^{-1} . Variety Vardan had the lowest protein content in seeds but produced significantly highest protein yield ha^{-1} owing to its highest grain yield during both years. Such differences have also been observed by several workers in their studies. Saran and De (2009), Kumar and Malik (2003), Singh (2009) reported variation in protein content in mustard seeds due to varieties.

1.2 REVIEW OF LITERATURE

The literature on the work proposed pertaining to the effect of nitrogen and sulphur nutrition on the growth, yield and quality of mustard varieties has been briefly reviewed here :

Effect of Varieties :

Gaur and Bansal (2013) showed that higher Chlorophyll and iron content of Pusa Bold leaves clearly indicated the effect of increased uptake of nitrogen under variety Pusa Bold compared to that of T-59 and Kranti. Similar results were reported by Gangasaran and Giri (2014) who found better response of Pusa Bold to nitrogen as compared to other mustard cultivars. Gaur et al. (2013) reported on the basis of two years experimentation that different cultivars significantly affected the fatty acid composition of mustard oil. Mustard variety T-59 recorded increase in erucic acid by 3.9 percent and a decrease in oleic acid content by 4.11 per cent of oil over that of variety RL-18. Singh et al. (2013) found that BR 40 was most susceptible to Zink deficiency whereas Pusa Bold was relatively tolerant, among six mustard varieties tested at Samastipur, Bihar. The relative susceptibility of the mustard varieties was rated as : RAUT-5-17 > BR 40 > BR 23 > Varuna > Kranti > Pusa Bold, on the basis of seed and oil yield response. Chandrakar et al (2013) reported that the seed yield of all the varieties of mustard showed similar yields with November 23 sowing over later dates.

1.3 OBJECTIVE OF THE STUDY :

- To study the comparative performance of mustard varieties under semi-arid conditions.
- To study the effect of nitrogen, sulphur and their interaction on the growth, yield and quality of mustard varieties.
- To work out the economic optimum doses of nitrogen and sulphur for mustard.

1.4 MATERIAL & METHOD

TECHNICAL PROGRAMME OF THE WORK

Details of Experiments :

Crop	:	Mustard (Brassica juncea L.)
Design	:	Split Plot Design
Replication	:	3

Treatments :

Main Plot :

(A) Varieties

V ₁	:	Vardan
V ₂	:	Jai Kisan (BIO 902)
V ₃	:	R H - 30

(B) Nitrogen levels (kg N/ha)

N ₀	:	0
N ₁	:	50
N ₂	:	100

Sub Plot :	Sulphur levels (kg S/ha)
S ₀	: 0
S ₁	: 25
S ₃	: 50

Treatment combinations :

$$3 \times 3 \times 3 = 27$$

Total Number of Plots :

$$= 27 \times 3$$

$$= 81$$

Plot Size :

$$\text{Gross} = 5.0 \text{ m} \times 3.15 \text{ m}$$

$$\text{Net} = 4.0 \text{ m} \times 1.35 \text{ m}$$

1.5 Effect of Nitrogen :

Repeseed and mustard need adequate supply of nitrogen for their growth. Its application enhances vegetative growth which is manifested through better plant height, branches and number of silique/plant which lead to higher seed yield.

Gangasaran and Giri (2018) reported that application of 80 kg N/ha, half at sowing and another half at 30 DAS resulted in significantly higher seed yield of irrigated mustard cv. Pusa Bold over other levels of nitrogen during Rabi 2015-16 and 2016-17. Aulakh et al. (2010) conducted field experiments for 3 years to test the effect of nitrogen on yellow mustard and black mustard on soils deficient in available nitrogen and reported that in all the 3 years seed yield of both mustard crops increased with increasing nitrogen levels. Sounda et al. (2009) observed that 90-100 kg N/ha was optimum for seed yield of irrigated mustard grown in sandy loam soil at Bichpuri, Agra (U.P.) reported that 90 kg N/ha benefited yield attributes and produced more amount of seed of irrigated mustard compared to other lower levels of nitrogen viz. 0, 30 and 60 kg N/ha.

1.6 Effect of Sulphur

Sulphur has long been recognised as essential for plant growth although much is to be learned about the function of this element. Sulphur is a constituent of the amino acid, methionine, cysteine and cystine. Plants that are sulphur deficient are characteristically small and spindly. The younger leaves are light green to yellowish.

Vaidya et al (2011) conducted trails at Udaipur in 2015-16 on mustard, grown on non-calcic brown soils and given 0-50 kg S/ha and found that average effect of sulphur was significant in increasing seed yield. Aulakh et al (2010) reported that sulphur fertilizer significantly increased the concentration and uptake (kg/ha) of sulphur in grains of yellow and black mustard. Application of 60 kg S/ha increased the oil content by 12.0 percent in yellow mustard.

Sawarkar et al. (2017) found that increasing the rates of applied sulphur 0-60 kg/ha increased the average oil content in mustard from 40.46 to 45.05 percent, decreased average protein content from 18.84 to 17.48 per cent, and also decreased S content from 0.28 to 0.26 per cent. Dubey and Khan (2013) concluded that 90 kg N and 30 kg S/ha would be the optimum doses for realizing the best growth of plant, seed yield, oil yield and N contents of mustard crop under irrigated condition of deep vertisol in Central Narmada Valley. Pathak and Tripathi (2009) reported from Kanpur that the seed yield of *B. juncea* was the highest with the application of 80 kg S/ha.

REFERENCES

- Aulakh, M.S., Pasricha, N.S. and Sahota, N.S. (2010). Yield nutrient concentration and quality of mustard crop as influenced by nitrogen and sulphur fertilizers J. Agri, Sci. 94(3) : 545-549.
- Chahal, R.S., Singh, B.P., Gupta, A.P. and Ram Kala (2011). Production potential of various crops under

- different levels of fertilizers and irrigation. *Indian J. Bot.* 25 : 258-261.
- Chandrakar, B.L. and Urkurkar, J.S. (2013). Performance of mustard (*B. juncea*) varieties to dates of sowing in rice fallow, *Indian J. Bot.* 38(1) : 143-144.
 - Chauhan, C.P.S., Singh, S.P., Pathak, D.S. and Bhudayal (2018). Nitrogen and phosphorus requirement of mustard crop under irrigation with saline water *Ann. Arid zones.* 27(3-4) : 293-296.
 - Dubey, O.P. and Khan, R.A. (2013). Effect of nitrogen and sulphur on dry matter, grain yield and nitrogen content at different growth stages of mustard (*B. juncea*) under irrigated vertisol. *Indian J. Agron.* 38(2) : 270-276.
 - Economic Survey. (2014). Annual Report. Economic division, Ministry of Finance, Govt. of India.
 - Gangasaran and Giri, G. (2015). Agronomic manipulations for increasing yields of edible oilseeds (In). *Oilseed Production Constraints and Opportunities.* (Eds.). H.C. Srivastava et al., oxford and IBH Pub. Co. Ltd., New Delhi. PP 285-292.
 - Gangasaran and Giri, G. (2017). Influence of irrigation and time of nitrogen application on mustard (*B. juncea*). *Indian J. Agron.* 33(2) : 154-158.
 - Gangasaran and Giri, G. (2014). Studies on comparative performance of mustard varieties under dryland conditions. *Indian J. Agron.* 29(3) : 398-401.
 - Gaur, B.L. and Bansal, N.K. (2013). Effect of varieties and mixtalol on leaf chlorophyll, ferrous content and nitrogen uptake of mustard (*B. juncea* (L) Czern and Coss). *J. Oilseeds Res.* 10(2) 1993 : 303-304.
 - Gaur, B.L., Verma, J. K. and Singh, H.G. (2013). Effect of nitrogen fertilization and plant population on fatty acid composition of select mustard (*B. juncea* L. Czern and Coss) cultivars, *J. Oilseeds Res.* 10(2) 1993 : 240-245.
 - Hearth, H.Y.M. and Ormrol, D.P. (2011). Temperature effect on the response to sulphur of barley, peas and rape. *Pl. Soil* 35 : 635-46.
 - Naik, M.S. and Das, N.B. (2004). Available S status of Indian soils by the *Aspergillus niger* method. *JISSS.* 12(2) : 151-156.
 - Pathak, R.K. and Tripathi, R.d. (2009). Effect of nitrogen and sulphur on yield and quality of Rai (*B. juncea*. L. var. Veruna). *Indian J. Agric. Chem.* 12(2) : 151-157.
 - Sawarkar, N.J., Shukla, S.S. and Tomar, P.S. (2007). Effect of sulphur fertilization on yield and quality of mustard. *J. Oilseed Res.* 4(2) : 227-229.
 - Singh, A.P., Sakal, R., Sinha, R.B. and Bhogal, N.S. (2013). Seed and Oil yield of mustard varieties as affected by Zn application in calcareous soil. *Ann, Agric. Res.* 14(4) : 457-462.
 - Sounda, G., Chatterjee, P. and Mandal, P.K. (2009). Effect of levels of nitrogen and moisture conservation practices on the yield and yield attributes of mustard at rainfed conditions. *Env. Ecol.* 7(1) : 186-188.
 - Tiwari, A.S. (2005). Rapeseed-mustard production in Madhya Pradesh, Contraints and Opportunities (Edited by Srivastava, H.C., Bhaskaran, S., Vatsya, B : and Menon, K.K.G.). New Delhi, India, Oxford and IBH Pub. Co. : 135-140.
 - Vaidya, P.K., Singh, M. and Nathwat, D.S. (2001). Response of mustard to the application of N, P, K and S. *Transactions of Indian Society of Desert Technology and University Centre of Desert Studies* 6(1) : 58-61.